

# **Inclusion criteria for Homeopathic treatment in**

## **Oncology**

### **LUNG CA-**

1. Primary lung cancer in cases with other co- morbidities making the patient not fit for chemo radiation.
2. Metastatic cases of lung CA in aged patients where chemo radiation have less chances.
3. In Lung CA after chemo radiation where reduction is not complete.

### **LIVER CA-**

1. Primary liver CA where TACE or RFA chances are less.
2. Metastatic CA where multiple liver lesions are there along with other secondaries.
3. To control AFP.

### **PANCREAS CA-**

1. In cases after whipples procedure.
2. In cases with upsurge of CA-19.9.
3. In cases where surgery/ Chemo radiation is not possible.

### **BREAST CA-**

1. Only in cases where chemo radiation/ Surgery is not possible.
2. Metastatic cases of CA breast especially with brain and bone secondaries to address sufferings .
3. After MRM with Axillary clearance in cases , to avoid adjuvant chemo radiation.

### **PROSTATE CA-**

1. In cases where skeletal metastasis in aged cases.
2. After TURPT to avoid Metastasis to bones.
3. Metastatic cases where chemo radiation is not possible.
4. To control PSA.

#### **RECTAL CA-**

1. In cases where colostomy is not possible .
2. In cases where adjuvant chemo radiation fails to reduce morbidities.
3. In primary cases in aged patients only.
4. In Metastatic cases to reduce metastatic foci after chemo radiation(liver/peritoneum).

#### **STOMACH CA-**

1. In cases of linitis plastica where surgery is not feasible.
2. In cases where chemo radiation is not possible due to anemia and emaciation.
3. After partial or complete gastrectomy to deal with metastasis to liver.
4. To regain physiological equilibrium in case under Feeding jejunostomy(FJ).

#### **ESOPHAGUS CA-**

1. Only in cases where surgery/ chemo/ radiation is contra indicated.
2. After chemo radiation or surgery to regain vitality.

#### **CERVICAL CA-**

1. In cases where surgery or radiation is not possible.
2. After chemo radiation to deal with metastasis.
3. After hysterectomy to avoid metastasis in aged cases.

#### **ORAL CA-**

1. Only in cases where medical condition contra indicative of conventional interventions.

#### **LYMPHOMA-**

1. In cases where chemotherapy is not possible.
2. In case of aged and multiple lymphadenopathy after relapse of scheduled dosage of chemotherapy.